



## Dissipation of Spirotetramat and Spirotetramat-Enol in Leaf and Fruit of Grape Variety Red Globe (*Vitis Vinifera* L.)

Norma J. Salazar-Lopez<sup>1</sup>, Maria I. Silveira- Gramont<sup>1</sup>, Fabiola G. Zuno-Floriano<sup>2</sup>, Guillermo Rodríguez-Olibarria<sup>1</sup>, Matt J. Hengel<sup>2</sup>, Ana I. Valenzuela-Quintanar<sup>3</sup>, Maria L. Aldana-Madrid<sup>1\*</sup>

<sup>1</sup>Departamento de Investigación y Posgrado en Alimentos, Universidad de Sonora, Hermosillo, Sonora, México.

<sup>2</sup>Department of Environmental Toxicology, University of California, Davis, One Shields Avenue Davis, CA. 95616, USA.

<sup>3</sup>Centro de Investigación en Alimentación y Desarrollo, A.C. Hermosillo, Sonora, México.

\*Corresponding author:

Maria L. Aldana-Madrid<sup>1</sup>

Phone: +526-622-592-208

### Abstract

The dissipation of spirotetramat (sp) and spirotetramat-enol (sp-enol) was determined in leaves and fruit of grapevine (*Vitis vinifera* L.) in a commercial vineyard. sp and sp-enol were extracted with acetonitrile:water, and quantification of residues was done on LC-DAD/MS. The LOQ of this method was found 0.002 mg kg<sup>-1</sup>, LOD being 0.001 mg kg<sup>-1</sup> for both. Recovery rates were in the range of 92 to 103% with coefficients of variation less than 9 and 6%, respectively at three levels (0.02, 0.2 and 0.6 mg kg<sup>-1</sup>). sp degradation in leaves followed first-order kinetics with R<sup>2</sup>=0.84 (p < 0.03). In fruit, only 35% of the original concentration of sp was detected at 5 days after application. These results suggest that residual levels of sp and sp-enol in grapes grown under the conditions of this study pose minimal risk to consumers, since residues did not exceed the limit established by the EPA.

**Keywords:** Spirotetramat; Spirotetramat-Enol; Grape; Dissipation; Grapevine.

### 1. Introduction

The application of agrochemicals to control crop pests may result in their presence in foods (Aldana et al., 2008a; Aldana et al., 2008b; Aldana et al., 2011) together with their physicochemical properties and environmental conditions under which they are applied (Babczynski and Hellpointner, 2008). Their residues may be of interest when products receive no processing before consumption, as in the case of table grapes (Oliva and Beard, 2002; Han et al., 2013).

Sp is an insecticide derived from the ambimobile insecticide tetramic acid aimed at the control of sucking pests in their juvenile stages (Nauen et al., 2008). It acts as an inhibitor of lipid biosynthesis, causing incomplete ecdysis, significant reduction in fecundity and fertility of insects and death (Fischer and Weiß, 2008; Kühnhold et al., 2008; Schöning, 2008). In experimental animals sp can cause eye irritation, skin sensitization, sperm abnormalities, hypospermia, tubular degeneration, and decreased testicular weight (FAO, 2008). Meanwhile, they may be the cause of contact dermatitis in humans (Funk, 2011). In toxicological studies in *Cerioda phniadubia*, it was reported that the mixture of an agricultural adjuvant (Destiny) and sp produced stronger toxicological effects (Chen and Stark, 2010). Even with its advantages and disadvantages, sp is a new alternative for the control of *Planococcus ficus*, which is a current problem in the cultivation of grapes on the coast of Sonora (Fu and del Real, 2009; Mansour et al., 2011). Accordingly, the objective of this study was to determine the dissipation of sp and sp-enol in leaves and fruit during cultivation of the grape variety (*Vitis vinifera* L.) Red Globe.

## 2. Materials and Methods

We used the standards of sp (cis-4-(ethoxycarbonyloxy)-8-methoxy-3-(2,5-xylyl)-1-azaspiro[4.5]dec-3-en-2-one) with 99.3% purity (Chem Service, West Chester, PA) and its enol metabolite (cis-3-(2,5-dimethylphenyl)-4-hydroxy-8-methoxy-1-azaspiro[4.5]dec-3-en-2-one) with 98.5% purity (Dr. Ehrenstorfer GmbH, Germany), HPLC grade acetonitrile (JT Baker, USA), reagent grade formic acid (88%) and SPE LC-18 cartridges (Supelco analytical No. 505471, USA). sp (Movento OD) at 15% was used in the field, where a dose of 600 cc per hectare was applied, equivalent to 90 g ha<sup>-1</sup> of active ingredient. The insecticide was applied by spraying on the foliage in a volume of 1500 L of water per hectare. During application, the temperature was 23.4°C, wind speed 2.1 m s<sup>-1</sup>, relative humidity 37.5% and solar radiation 0.331 kW m<sup>-2</sup>. The samples were collected in San Luis Vineyard located at Km 19.5 of North 12th Street in Costa de Hermosillo, Sonora, Mexico at 29° 00' 53'' N, 111° 29' 16'' W (Datum WGS84) at an altitude of 78 m. The cultivation area consisted of 106 rows of plants separated by four meters, each row with 182 vines separated by 1.4 m. A drip irrigation system was used, set approximately 50 cm from the ground. Within the vineyard, five points were randomly selected, where plots were formed which in turn were made up of three plants located in the same row, for collection of samples of leaves and fruits at six time points; the day before the application of insecticide and on days 1, 5, 10, 15 and 48 after application. About 300 g of each sample were collected, placed in plastic bags and stored at -20°C on a laboratory freezer until analysis.

### Detection and Quantification of Sp and Sp-Enol by LC-MS

The instrumentation used was an Agilent Technologies 1100 Series LC-MS chromatograph coupled to a simple quadrupole mass spectrometer model VL and DAD, equipped with a quaternary pump (1310a) and G1313A autosampler. Separation was carried out with an XBridge column, 3.5 µm and 3.0 x 150 mm, and XBridge C18 precolumn (Waters), 3.5 µm and 3.0 x 20 mm. Data were collected using Rev B.03.02 Chemstation software. The analysis of sp and sp-enol was according to Schöning (2008). The precursor ions and quantification of sp and sp-enol were 374→216, 302 m/z and 302→216, 270 m/z, respectively. The retention times for sp and sp-enol were 9.17 and 7.6 minutes, respectively. Satisfactory linearities were observed for sp and sp-enol ( $R^2 \geq 0.9926$ ) for both compounds.

### Statistical Analysis

We used a completely randomized design, with six repeated observations on days 0, 1, 5, 10, 15 and 48, after application of sp, and five replicates. Response variables were the amounts of sp and sp-enol on fruits and leaves. Descriptive statistics were carried out for each time in leaves and fruit. Analysis of variance with repeated measures was performed to determine: a) persistence [difference between initial time (before) and time with maximum concentration detected] and b) assessment of dissipation profiles of the insecticide and its metabolite in the plant.

## 3. Results and Discussion

Spirotetramat recovery percentages obtained for levels of 0.02, 0.2 and 0.6 mg kg<sup>-1</sup> are show on **Table 1**. The results were similar to those reported by Singh et al. (2013) in okra, eggplant and peppers (82-97%) and Schöning (2008) in tomato, avocado, citrus, potato and hops (84-104%), and higher than those reported by Mohapatra et al. (2012) in mango and cabbage (sp and sp-enol respectively 73-93 and 84-116% in mango; sp and sp-enol respectively 74-91 and 87-109% in cabbage).

**Table 1: Spirotetramat Recovery Percentages Obtained for Levels of Compounds**

Compound	Compound level (mg kg <sup>-1</sup> )	Leaves (%)	Immature fruit (%)	Ripe fruit (%)
Spirotetramat	0.02	99 ± 5	101 ± 8	98 ± 3
	0.2	98 ± 3	96 ± 1	95 ± 4
	0.6	99 ± 1	100 ± 2	93 ± 4
Spirotetramat-enol	0.02	99 ± 6	99 ± 2	92 ± 3
	0.2	103 ± 2	98 ± 1	97 ± 4
	0.6	99 ± 2	103 ± 9	93 ± 5


n=5

The dissipation of sp in leaves was fitted to a first-order kinetics model with  $R^2 = 0.84$  ( $p < 0.03$ ), the initial concentration found a day after the application was  $0.14 \text{ mg kg}^{-1}$ ; after 5 days, there was a dissipation of 13%, and after 48 days, this increased up to 79%, resulting in a concentration of  $0.03 \text{ mg kg}^{-1}$ . The half-life of the insecticide in leaves was 9 days with a 95% confidence interval of (7-11). Dissipation data were similar to those previously reported by Maclachlan and Hamilton (2010) for the degradation of sp in apple. sp degradation in fruit was 65% during the first 5 days after application, giving a concentration of  $0.08 \text{ mg kg}^{-1}$ ; at 48 days after the application, a residual concentration of  $0.02 \text{ mg kg}^{-1}$  was detected. Vemuri et al. (2014) reported a dissipation rate less than that observed in this study, possibly because of the combined application of imidacloprid and sp on grapevines. However, in both studies, concentrations below the maximum residue limit (MRL) established by the EPA (2011) ( $1.3 \text{ mg kg}^{-1}$ ) are reported. In this study, the half-life of sp in fruit was  $5 \pm 2$  days. These results differed from those reported by Cahil et al. (2015), who obtained a half-life of 1.91 days for sp in the fruit of chili pepper; the shorter half-life could be due to the high relative humidity of the crop (48-100%). Residual levels of sp were higher in leaves compared to fruit (Fig. 1A). This behavior was previously reported by Sur (2008), who evaluated the concentration of sp in the fruit and leaves of apple, as well as in the tuber and leaves of potato, finding a higher concentration in the leaves in both cases. This was possibly due to the greater density of leaves than fruit of the plant, and thus leaves may be exposed to a larger amount of insecticide. The results for sp-enol in leaves and fruit at 5 days after application showed 70 and 80% dissipation, respectively, of the residue in relation to the concentration at day 1 ( $0.03$  and  $0.04 \text{ mg kg}^{-1}$ ). After day 5, residual levels slowly decreased to  $0.005$  and  $0.008 \text{ mg kg}^{-1}$  in leaves and fruit, respectively. It was clear that this metabolite showed different degradation kinetics as sp, but it was not possible to determine the dissipation kinetics of sp-enol in leaves and fruit, since the data showed no significant differences after day 5 (Fig. 1B). It would be interesting in future studies to take samples daily, to elucidate its degradation kinetics.

The sp dissipation model (first-order kinetics) coincides with that reported for the ketoenols spirodiclofen and spiromesifen (Sharma et al., 2006), and likewise, the sp-enol metabolite was detected at 12 hours after application of the pesticide. In the present investigation, sp-enol metabolite was first detected at 24 hours, since no samples were taken on the same day the insecticide was applied.

Sharma et al. (2007a) in their investigation of the dissipation kinetics of spiromesifen in tea (*Camellia sinensis*) grown under tropical conditions, showed that 99% of the compound was degraded between 33 and 57 days after application. Also, in chili pepper and cotton, 99% degradation of spiromesifen occurred between 14.5 and 16.8 days by first-order kinetics (Sharma et al., 2007b).

The results obtained in this study suggest that under the agricultural practices used in growing Red Globe table grapes in the state of Sonora, Mexico, residual levels of sp and its enol metabolite pose minimal risk to consumers, since they are below the MRLs established by the EPA. However, it would be interesting to conduct future studies with daily sampling after application of the insecticide and to determine the dissipation of sp and the formation and dissipation of its degradation products such as enol-glucoside, ketohydroxy and monohydroxy metabolites.



**Fig 1: Dissipation of sp (A) and sp-enol (B) in leaves and fruit of grapevine with respect to time**

## Acknowledgements

This study has been carried out with financial support from the University of Sonora (Mexico) and Consejo Nacional de Ciencia y Tecnología (CONACyT) scholarship 377167. The authors sincerely thank Rubén Romo (San Luis vineyard) for providing samples. Riza Punongbayan and Susana-Alejandra Palma-Durán for technical support.

## 4. References

- [1] Ahmed AR, Tarek MM, Rady AR, Mohamed YH (2009). Dissipation of profenofos, imidacloprid and penconazole in tomato fruits and products. *Bull Environ Contam Toxicol* 83:812-817.
- [2] Aldana ML, García MM, Rodríguez OG, Silveira GM, Valenzuela QA. (2008a). Determinación de insecticidas organofosforados en nopal fresco y deshidratado. *Rev Fitotec Mex* 31(2):133-139 Available <http://redalyc.uaemex.mx/redalyc/src/inicio/ArtPdfRed.jsp?iCve=61031206>.
- [3] Aldana ML, Valdez S, Vargas ND, Salazar NJ, Silveira MI, Loarca FG, Rodríguez G, Wong FJ, Borboa J, Burgos A (2008b). Insecticide residues in stored grains in Sonora, Mexico: quantification and toxicity testing. *Bull Environ Contam Toxicol* 80:93-96.
- [4] Aldana ML, Valenzuela AI, Silveira MI, Rodríguez G, Grajeda P, Zuno FG, Miller MG (2011). Residual pyrethroids in fresh horticultural products in Sonora, Mexico. *Bull Environ Contam Toxicol* 87(4):436-439. DOI 10.1007/s00128-011-0391-z.
- [5] Alsayeda H, Pascal-Lorber S, Nallanthigal C, Debrauwer L, Laurent F (2008). Transfer of the insecticide [14C] imidacloprid from soil to tomato plants. *Environ Chem Lett* 6:229–234.
- [6] B /abczinski P, and Hellpointner E (2008). Environmental fate of spirotetramat (Movento). *Bayer Crop Sci J* 61(2):181-201.
- [7] Chalil GS, Mandal K, Sahoo SK (2015). Risk assessment of mixture formulation of spirotetramat and imidacloprid in chilli fruits. *Environ Monit Assess* 187:4105(1-8).
- [8] Chen XD, and Stark JD (2010) Individual -and population- level toxicity of the insecticide, spirotetramat and the agricultural adjuvant, Destiny to the Cladoceran, *Ceriodaphniadubia*. *Ecotoxicology* 19:1124–1129.
- [9] EPA United States Environmental Protection Agency (EPA) (2011) <http://www.gpo.gov/fdsys/pkg/FR-2011-05-18/html/2011-11937.htm> Accessed 27 February 2015.
- [10] FAO (2008) Pesticide residues in food 2008 joint FAO/WHO meeting on pesticide residues. <http://www.fao.org/docrep/011/i0450e/i0450e00.htm> Accessed 27 February 2015.
- [11] Fisher R, and Weiß HC (2008). Spirotetramat (Movento®) – Discovery, synthesis and physicochemical properties. *Bayer Crop Sci J* 61(2):127-140.
- [12] Fu AA, and del Real AA (2009). Guía para el control de piojo harinoso de la vid. Folleto Técnico N° 38. Centro de investigación Regional del Noroeste Campo Experimental Costa de Hermosillo. Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias.
- [13] Funk S (2011). SPIROTETRAMAT. FAO and Environmental Protection Agency, USA. [http://www.fao.org/fileadmin/templates/agphome/documents/Pests\\_Pesticides/JMPR/Evaluation08/Spirotetramat.pdf](http://www.fao.org/fileadmin/templates/agphome/documents/Pests_Pesticides/JMPR/Evaluation08/Spirotetramat.pdf) Accessed 11 February 2015.
- [14] García MA (2001). Influencia de los residuos de fungicidas en la cinética fermentativa y calidad de los vinos blancos de la D.O. Jumilla. Universidad de Murcia, España.
- [15] Han Y, Xu J, Dong F, Li W, Liu X, Li Y, Kong Z, Zhu Y, Liu N, Zheng Y (2013). The fate of Spirotetramat and its metabolite Spirotetramat-enol in apple samples during apple cider processing. *Food Chem* 34:283-290.
- [16] Kühnhold J, Klueken AM, de Maeyer L, Waetermeulen XV, Brück E, Elbert A (2008). Movento®, an innovative solution for sucking insect pest control in agriculture: field performance in fruits and vegetables. *Bayer Crop Sci J* 61(2):279-306.
- [17] Maclachlan DJ, and Hamilton D (2010). A new tool for the evaluation of crop residue trial data (day-zero-plus decline). *Food Addit Contam* 27(3):347-364.
- [18] Mansour R, Suma P, Mazzeo G, Lebdi KG, Russo A (2011). Evaluating side effects of newer insecticides on the vine mealybug parasitoid *Anagyrus* sp. near *pseudococci*, with implications for integrated pest management in vineyards. *Phytoparasitica* 39:369-376.

- [19] Mohapatra S, Deepa M, Jagadish GK (2012). An efficient analytical method for analysis of spirotetramat and its metabolite spirotetramat-enol by HPLC. *Bull Environ Contam Toxicol* 88:24-128.
- [20] Murcia AM, and Stashenko E (2008). Determinación de plaguicidas organofosforados en vegetales producidos en Colombia. *Agro Sur* 36(2):71-81.
- [21] Nauen R, Reckmann U, Thomzik J, Thielert W (2008). Biological profile of spirotetramat (Movento®) – a new two-way systemic (ambimobile) insecticide against sucking pest species. *Bayer Crop Sci J* 61(2):245-277.
- [22] Oliva J, and Barba A (2002) Elaboración de vino en presencia de residuos de plaguicidas. Alteraciones, control y desaparición. *Tecnología del vino* (pp. 45-49). Available <http://www.alcion.es/Download/ArticulosPDF/tv/03/TVMA3DOC.pdf>.
- [23] Piñero M, Izquierdo P, Cagnasso MA, García A (2007). Residuos de plaguicidas organoclorados en 4 tipos de aceites vegetales. *Arch Latin Nut (ALAN)* 57(4):397-401. Available <http://www.scielo.org.ve/pdf/alan/v57n4/art13.pdf>.
- [24] Schöning R (2008). Residue analytical method for the determination of residues of spirotetramat and its metabolites in and on plant material by HPLC-MS/MS. *Bayer Crop Sci J* 61:417-454.
- [25] Sharma KK, Dubey JK, Deka SC, Chandrasekaran S, Kalpana, Gupta P, Kumar A, Vandana, Jaya Devi M, Singh B, Baruah AA, Kennedy JS, Sharma ID, Patyal SK, Nath A (2007a). Dissipation kinetics of spiromesifen on tea (*Camellia sinensis*) under tropical conditions. *Chemosphere* 68(4):790-796.
- [26] Sharma KK, Cherukuri SR, Dubey JK, Patyal SK, Parihar NS, Battu RS, Sharma V, Gupta P, Kumar A, Kalpana, Jaya Devi M, Singh B, Sharma ID, Nath A, Gour TB (2007b). Persistence and dissipation kinetics of spiromesifen in Chili and cotton. *Environ Monit Assess* 132:25-31.
- [27] Sharma KK, Dubey JK, Mukherjee I, Parihar NS, Battu RS, Singh B, Kumar A, Gupta P, Kalpana B, Sharma ID, Nath A (2006). Residual behavior and risk assessment of spiromesifen (Oberon 240 SC) on eggplant (*Palampurum melonongena* L) in India: A multilocational study. *Bull Environ Contam Toxicol* 76(5): 760-765.
- [28] Singh B, Kousik M, Sanjay S, Urvashi B, Raminderjit B (2013). Development and validation of an HPLC method for determination of spirotetramat and spirotetramat cis enol in various vegetables and soil. *J AOAC Int* 96(3):670-675.
- [29] Sur, R. (2008). Metabolism of spirotetramat (Movento®) in plants. *Bayer Crop Sci J* 61(2): 227-243.
- [30] Vemuri S, Rao CR, Swarupa S (2014) Dissipation of spirotetramat and imidacloprid in grapes and soil. *J Mult Eng Sci Tech (JMEST)* 1(5):319-324.